

ProSci Incorporated 12170 Flint Place Poway, CA 92064, USA prosci-inc.com P: +1 (888) 513-9525 Local: +1 (858) 513-2638 Fax: +1 (858) 513-2692

PD-L2 Antibody


CATALOG NUMBER: 4063

Specifications

Host Species	Rabbit
Species Reactivity	Human, Mouse
Homology	Predicted species reactivity based on immunogen sequence: rat (94%).
Immunogen	Anti-PD-L2 antibody (4063) was raised against a peptide corresponding to 16 amino acids near the center of human PD-L2. The immunogen is located within amino acids 140-190 of PD-L2.
Conjugate	Unconjugated
Tested Applications	ELISA, IF, IHC-P, WB
User Note	Optimal dilutions for each application to be determined by the researcher.
Predicted Molecular Weight	Predicted: 31kD Observed: 29 kD


Properties

Purification	PD-L2 Antibody is affinity chromatography purified via peptide column.
Clonality	Polyclonal
Isotype	IgG
Physical State	Liquid
Buffer	PD-L2 Antibody is supplied in PBS containing 0.02% sodium azide.
Concentration	1 mg/mL
Storage Conditions	PD-L2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.




PD-L2 Antibody 1

Figure 1 Western Blot Validation in Human Raji Cell Lysate Loading: 15 μ g of lysates per lane.Antibodies: PD-L2 4063 (A: 0.5 μ g/mL and B: 1 μ g/mL), 1h incubation at RT in 5% NFD...




PD-L2 Antibody 3

Figure 3 KO Validation in HeLa Cells Loading: 15 μ g of HeLa WT cell lysates or PD-L2 KO cell lysates. Antibodies: PD-L2, 4063 (4 μ g/mL) and beta-actin 3779 (1 μ g/mL), 1 h incu...



PD-L2 Antibody 2

Figure 2 Independent Antibody Validation (IAV) via Protein Expression Profile in Human and Mouse Cell Lines Loading: 15 μ g of lysates per lane. Antibodies: PD-L2, 4063 (4 μ g/mL),...



PD-L2 Antibody 4

Figure 3 KO Validation in HeLa Cells Loading: 15 μ g of HeLa WT cell lysates or PD-L2 KO cell lysates. Antibodies: PD-L2, 4063 (4 μ g/mL) and beta-actin 3779 (1 μ g/mL), 1 h incu...

Disclaimer

Disclaimer

Optimal dilutions/concentrations should be determined by the end user. The information provided is a guideline for product use. This product is for research use only.

For research use only. For additional information, visit ProSci's Terms and Conditions Page.